

Supplementary Table 2. Changes in systemic parameters in study participants relative to changes in central subfield retinal thickness

Patient	Drug Adherence	Duration of Type II Diabetes (years)	HbA1c (mg/dL)			Creatinine (mg/dL)			Blood Pressure (mmHg)				Visual Acuity (letters)			OCT Central Subfield Thickness (microns)			
			BL	Month 6	Change	BL	Month 6	Change	BL	Month 6	Change in systolic	Change in diastolic	BL	Month 6	Change	BL	Month 6	% Change	
1	98.9%	3	6.7	6.6	-0.1	0.88	0.88	0	145/80	121/74	-24	-6	Study Eye	65	85	20	298	240	-19.5%
													Fellow Eye	88	88	0	289	257	-11.1%
2	98.7%	18	9.9	9.3	-0.6	1.24	0.92	-0.32	119/81	138/81	19	0	Study Eye	85	81	-4	257	272	5.8%
													Fellow Eye	40	30	-10	139	523	276.3%
3	100.3%	50	7.8	8	0.2	0.91	0.88	-0.03	110/64	124/68	14	4	Study Eye	81	71	-10	352	302	-14.2%
													Fellow Eye	83	85	2	267	250	-6.4%
4	96.2%	31	7.4	7.5	0.1	1.35	1.46	0.11	111/51	126/55	15	4	Study Eye	65	62	-3	331	377	13.9%
													Fellow Eye	80	81	1	292	315	7.9%
5	94.1%	6	8	8.4	0.4	0.67	0.74	0.07	125/70	127/70	2	0	Study Eye	80	80	0	438	361	-17.6%
													Fellow Eye	79	75	-4	356	466	30.9%
Mean	97.6%	21.6			0			-0.034			5.2	0.4				-0.8		26.6%	
SD	2.5%	19.3			0.4			0.2			17.5	4.1				8.4		89.2%	